

# Convex Functions, Transformations

Huanle Xu \*

April 9, 2018

## 1 Convex Optimization Problem

All convex optimization problems have the form of:

$$\begin{aligned} \min_x & f(x) \\ \text{s.t. } & g_i(x) \leq 0, \quad \forall i = 1, 2, \dots, m \\ & x \in X, \end{aligned}$$

where  $f(x)$  and  $g_i(x)$  are convex functions and  $X \in \mathbb{R}^n$  is a convex set.

## 2 Convex functions

Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$ , we say that  $f$  is convex if

$$f(\alpha x_1 + (1 - \alpha)x_2) \leq \alpha f(x_1) + (1 - \alpha)f(x_2), \quad (1)$$

for all  $x_1, x_2 \in \mathbb{R}^n$  and  $\alpha \in [0, 1]$ . We say that  $f$  is concave if  $-f$  is convex.

**Proposition 1.** *Consider the optimization problem*

$$\begin{aligned} \min_x & f(x) \\ \text{s.t. } & x \in S, \end{aligned}$$

where  $S \in \mathbb{R}^n$  is a convex set and  $f$  is a convex function. Then, any local minimizer is also a global minimizer.

### 2.1 Convexity-Preserving Transformations

The following hold:

- (Non-Negative Combinations) Let  $f_1, \dots, f_m$  be convex functions, and let  $\alpha_1, \dots, \alpha_m \geq 0$ . Then, the function  $\sum_{i=1}^m \alpha_i f_i$  is also convex.
- (Pointwise Supremum) Let  $\{f_i\}_{i \in I}$  be an arbitrary family of convex functions on  $\mathbb{R}^n$ . Then, the pointwise supremum  $f = \sup_{i \in I} f_i$  is also convex.
- (Composition with an Increasing Convex Function) Let  $f$  be a convex function, and let  $g : \mathbb{R} \rightarrow \mathbb{R}$  be an increasing convex function. Then, the function  $g(f(x))$  is convex on  $\mathbb{R}^n$ .

---

\*Huanle Xu is with the College of Computer Science and Technology, Dongguan University of Technology. E-mail: xuhl@dgut.edu.cn.

## 2.2 Differentiable Convex Functions

When  $f$  is a differentiable function, we can characterize its convexity via its gradient.

**Theorem 1.** *Let  $f : \Omega \rightarrow \mathbb{R}$  be a differentiable function on the open set  $\Omega \in \mathbb{R}^n$ , and let  $S \subset \Omega$  be convex. Then,  $f$  is convex on  $S$  iff*

$$f(x_1) \geq f(x_2) + (\nabla f(x_2))^T(x_1 - x_2), \quad (2)$$

for all  $x_1, x_2 \in S$ .

**Theorem 2.** *Let  $f : S \rightarrow \mathbb{R}$  be a twice continuously differentiable function on the open convex set  $S \subset \mathbb{R}^n$ . Then,  $f$  is convex on  $S$  iff  $\nabla^2 f(\bar{x})$  is positive semidefinite for all  $\bar{x} \in S$ .*

## 3 Some Useful Inequalities

Let us begin with Jensen's inequality, which can be viewed as a generalization of (1).

**Proposition 2. (Jensen's Inequality)** *Let  $f$  be a convex function. Then, for any  $x_1, x_2, \dots, x_k \in \text{dom}(f)$  and  $\alpha_1, \alpha_2, \dots, \alpha_k \in [0, 1]$  such that  $\sum_{i=1}^k \alpha_k = 1$ , we have*

$$f\left(\sum_{i=1}^k \alpha_i x_i\right) \leq \sum_{i=1}^k \alpha_i f(x_i). \quad (3)$$

**Proposition 3.** *For all  $x_1, x_2, \dots, x_n \in \mathbb{R}_+$ , the following holds:*

$$\left(\prod_{i=1}^n x_i\right)^{1/n} \leq \frac{1}{n} \sum_{i=1}^n x_i. \quad (4)$$

## 4 Examples of Convex Functions

1. Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  be given by  $f(x) = \log\left(\sum_{i=1}^n \exp(x_i)\right)$ . We compute

$$\frac{\partial^2 f}{\partial x_i \partial x_j} = \begin{cases} \frac{\exp(x_i)}{\sum_{i=1}^n \exp(x_i)} - \frac{\exp(2x_i)}{(\sum_{i=1}^n \exp(x_i))^2} & \text{if } i = j, \\ -\frac{\exp(x_i + x_j)}{(\sum_{i=1}^n \exp(x_i))^2} & \text{if } i \neq j. \end{cases}$$

2. Let  $f : \mathbb{R}_+^n \rightarrow \mathbb{R}$  be given by  $f(x) = \left(\prod_{i=1}^n x_i\right)^{1/n}$ . We compute

$$\frac{\partial^2 f}{\partial x_i \partial x_j} = \begin{cases} -(n-1) \frac{(\prod_{i=1}^n x_i)^{1/n}}{n^2 x_i^2} & \text{if } i = j, \\ \frac{(\prod_{i=1}^n x_i)^{1/n}}{n^2 x_i x_j} & \text{if } i \neq j. \end{cases}$$